

# Symbolic solving of partial differential equations systems and compatibility conditions.

Natasha Malaschonok

namalaschonok@gmail.com

Tambov State University,

Internatsionalnaya 33, 392622 Tambov, Russia

1

**Abstract.** There is produced an algorithm for symbolic solving systems of partial differential equations by means of multivariate Laplace-Carson transform. There is considered a system of  $K$  equations with  $M$  as the greatest order of partial derivatives and right hand parts of a special type. Initial conditions are input. As a result of Laplace-Carson transform of the system according to initial condition we obtain an algebraic system of equations. A method to obtain compatibility conditions is discussed.

Laplace-Carson transform, systems of partial differential equations, symbolic solving, compatibility conditions

## 1 Introduction

The Laplace transform has been useful in various problems of differential equations theory, including problems of partial equations (for example [?] [?], [?],[?], [?],[?],[?]). It serves as a base for an operation calculus, that is used in such applications. We must mention, that kinds of operation calculus exist, such as Mikusinski-type operational calculus, (see for example [?],[?], [?], [?]), or an approach of M.Gutterman (for example [?]), close to ideas of J.Mikusinski, which use the convolution quotient, without referring to the Laplace transform. Our method permits not to reduce PDE equations to ordinary ones, but to obtain directly the algebraic system of equations for further activities with it.

On the other hand, there are many ways to use computer algebra systems for numerical or symbolic solving of PDE systems, for example the well known usage of MAPLE for characteristics method that permits to simplify equations in many cases (for instance [?],[?]).

The method, produced in this paper, is now of rising actuality because of increasing relevance of symbolic computations nowadays. It reduces a system of partial differential equations with initial conditions to an algebraic linear system with polynomial coefficients which already encloses in symbolic way the initial conditions requirements. An important advantage of the method is the

---

<sup>1</sup> Supported by RFBR, No.12-07-00755-a, RusMES No.1.345.2011

establishment of compatibility conditions in symbolic way for many types of PDE equations and systems of PDE equations. A type of PDE equations is not of great importance as well as their order or a size of a system.

We produce an algorithm for symbolic solving of systems of linear partial differential equations by means of multivariate Laplace–Carson transform. Considered are the systems of arbitrary number  $K$  of unknown functions and equations of arbitrary order  $M$  of derivatives. The method allows not to reduce to canonical form the problem at initial stage, it reduces it to solving a linear algebraic system with polynomial coefficients where efficient methods were developed (for example [?, ?, ?, ?]). A parallel algorithm may be constructed on the base of the produced one. A way of parallelization is briefly discussed. So large systems of linear PDE may be solved in real time.

## 2 Problem statement

Denote  $\tilde{m} = (m_1, \dots, m_n)$ . Consider a system

$$\sum_{k=1}^K \sum_{m=0}^M \sum_{\tilde{m}} a_{mk}^j \frac{\partial^m}{\partial^{m_1} x_1 \dots \partial^{m_n} x_n} u_k(x) = f_j(x), \quad (1)$$

where  $j = 1, \dots, K$ ,  $u_k(x)$ ,  $k = 1, \dots, K$ , — are unknown functions of  $x = (x_1, \dots, x_n) \in \mathbf{R}_+^n$ ,  $f_j \in S$ ,  $a_{mk}^j$  are real numbers,  $m$  is the order of a derivative, and  $k$  — the number of an unknown function. Here and further summing by  $\tilde{m} = (m_1, \dots, m_n)$  is executed for  $m_1 + \dots + m_n = m$ .

Functions  $f_j(x)$  of the right-hand part of the system are in general composite. For each  $j$  consider  $\mathbf{R}_+^n$  divided into parts  $D_j^i = \{x : x_{\nu,j}^{i_\nu} < x_\nu < x_{\nu,j}^{i_\nu+1}, \nu = 1, \dots, n\}$ ,  $i_\nu = 1, \dots, I_{\nu,j}$ ,  $x_{i_1} = 0, x_{I_{\nu,j}+1} = \infty$ .

$$f_j(x) = f_j^i(x), \quad x \in D_j^i \quad i = (i_1, \dots, i_n).$$

Denote the set of  $i$  for which  $i_\nu = 1, \dots, I_{\nu,j}$  by  $\mathbf{I}_j$ . As usually,  $x - x_i = (x_1 - x_{i_1}, \dots, x_n - x_{i_n})$ .  $H(x) = H(x_1, \dots, x_n)$  is the multivariate Heaviside function which equals 1 only if  $x_\nu \geq 0$  for all  $\nu$ , and zero otherwise. For  $l = (l_1, \dots, l_n)$  we denote  $x^l = \prod_{\nu=1}^n x_\nu^{l_\nu}$ ,  $\langle b_{js}^i, x \rangle = \sum_{\nu=1}^n b_{js}^{i_\nu} x_\nu$ . Using Heaviside function, composite functions  $f_j(x)$  may be written as follows

$$f_j(x) = \sum_{i \in \mathbf{I}_j} \left[ \widetilde{f_j^i}(x - x_i) H(x - x_i) - \widetilde{f_j^i}(x - x_{i+1}) H(x - x_{i+1}) \right], \quad (2)$$

where  $\widetilde{f_j^{I_j+1}}(x - x_{I_j+1}) = 0$ , and

$$f_j^i(x) = \sum_{s=1}^{S_j^i} P_{js}^i(x) e^{\langle b_{js}^i, x \rangle}, \quad i = 1, \dots, I_j, \quad j = 1, \dots, k,$$

$P_{js}^i(x) = \sum_{l=0}^{L_{js}^i} c_{sl}^{ji} x^l$  is a polynomial,  $L_{js}^i = (L_{js}^{i1}, \dots, L_{js}^{in})$ . For a polynomial  $P(x)$  we denote  $\tilde{P}(x - x_i)$  its expansion in powers of  $(x_1 - x_{i1}), \dots, (x_n - x_{in})$ , and for  $f(x) = P(x)e^{\langle a, x \rangle}$  we denote  $\tilde{f}(x - x_i) = \tilde{P}(x - x_i)e^{\langle a, x_i \rangle} e^{\langle a, x - x_i \rangle}$ .

Denote by  $\mathbf{A}$  a class of functions which are reducible to the form (2).

We solve a problem with initial conditions for each variable (not separating spatial or time data, it is defined by concrete problem). Introduce notations for them.

Denote by  $\Gamma^\tau$  a set of vectors  $\gamma = (\gamma_1, \dots, \gamma_n)$  such that  $\gamma_\tau = 1$ ,  $\gamma_i = 0$ , if  $i < \tau$ , and  $\gamma_i$  equals 0 or 1 in all possible combinations for  $i > \tau$ . The number of elements in  $\Gamma^\tau$  equals  $2^{\tau-1}$ .

Denote  $\beta = (\beta_1, \dots, \beta_n)$ ,  $\beta_i = 0, \dots, m_i$ , a set of indexes such that the derivative of  $u_k(x)$  of the order  $\beta_i$  with respect to the variables with numbers  $i$  equals  $u_{\beta, \gamma}^k(x^{(\gamma)})$  at the point  $x = x^\gamma$  with zeros at the positions  $\mu$  for which the coordinates  $\gamma_\mu$  of  $\gamma$  equal 1 (for example, if zeros stand only at the places with the numbers 1, 2, 3, then  $\gamma = (1, 1, 1, 0, \dots, 0)$ ). We take functions  $u_{\beta, \gamma}^k(x^{(\gamma)})$  also from the class  $\mathbf{A}$ .

Further we shall introduce a notion of compatible initial conditions and consider a method to obtain compatible initial conditions. The system (1) is to be solved under such conditions.

### 3 Laplace–Carson transform

Consider the space  $S$  of functions  $f(x)$ ,  $x = (x_1, \dots, x_n) \in \mathbf{R}_+^n$ ,  $\mathbf{R}_+^n = \{x : x_i \geq 0, i = 1, \dots, n\}$ , for which  $\mathcal{M} > 0$ ,  $a = (a_1, \dots, a_n) \in \mathbf{R}_+^n$ ,  $a_i > 0$ ,  $i = 1, \dots, n$ , exist such that for all  $x \in \mathbf{R}_+^n$  the following is true:  $|f(x)| \leq \mathcal{M}e^{\langle a, x \rangle}$ . Evidently,  $\mathbf{A} \subset S$ .

On the space  $S$  the Laplace–Carson transform (LC) is defined as follows:

$$LC : f(x) \mapsto F(p) = p^1 \int_0^\infty e^{-\langle p, x \rangle} f(x) dx,$$

$p = (p_1, \dots, p_n)$ ,  $p^1 = p_1 \dots p_n$ ,  $dx = dx_1 \dots dx_n$ . Integration is multivariate. The Laplace–Carson transform is more convenient than Laplace transform, as it transfers the Heaviside function into the unit, and it rather simplifies calculations.

The Laplace–Carson image of  $(x - x_i)^l e^{\langle \alpha, x - x_i \rangle} H(x - x_i)$  is

$$p^1 \prod_{\nu=1}^n \frac{l_\nu!}{(p_\nu - \alpha_\nu)^{l_\nu+1}} e^{-\langle p, x_i \rangle}, \quad \alpha = (\alpha_1, \dots, \alpha_n). \quad (3)$$

That is why LC is performed symbolically on the class  $\mathbf{A}$ .

Let  $LC : u_k \mapsto U_k, u_{\beta, \gamma}^k(x^{(\gamma)}) \mapsto U_{\beta, \gamma}^k(p^{(\gamma)}), f_j \mapsto F_j$ , the notation  $p^{(\gamma)}$  is correspondent to the notation  $x^{(\gamma)}$ . Denote by  $\|\gamma\|$  the “length” of  $\gamma$  — the number of units in  $\gamma$ .

The LC of the left-hand side of the system (1) is being written formally. Write at first the image of a derivative;

$$LC : \frac{\partial^m}{\partial^{m_1} x_1 \dots \partial^{m_n} x_n} u_k(x) \mapsto p^m U_k(p) + \sum_{\nu=1}^n \sum_{\beta_\nu=0}^{m_\nu} \sum_{\gamma \in \Gamma^\nu} (-1)^{\|\gamma\|} p_1^{m_1 - \beta_1 - \gamma_1} \dots p_n^{m_n - \beta_n - \gamma_n} U_{\beta, \gamma}^k(p^{(\gamma)}).$$

Denote

$$\Phi_{mk}^j = \sum_{\tilde{m}} a_{mk}^j \sum_{\nu=1}^n \sum_{\beta_\nu=0}^{m_\nu} \sum_{\gamma \in \Gamma^\nu} (-1)^{\|\gamma\|} p_1^{m_1 - \beta_1 - \gamma_1} \dots p_n^{m_n - \beta_n - \gamma_n} U_{\beta, \gamma}^k(p^{(\gamma)}).$$

As a result of Laplace–Carson transform of the system (1) according to initial conditions we obtain the algebraic system relative to  $U_k$ :

$$\sum_{k=1}^K \sum_{m=0}^M \sum_{\tilde{m}} a_{mk}^j p^{\tilde{m}} U_k(p) = F_j - \sum_{k=1}^K \sum_{m=0}^M \Phi_{mk}^j, j = 1, \dots, K. \quad (4)$$

## 4 Solution of algebraic system

Efficient methods of solving such systems are developed according to size or density of the algebraic system (for example [?, ?, ?, ?]).

Denote by  $D$  the determinant of the system (4), by  $D_k$  the minor that is obtained by replacing the  $k$ -th column by the column of the right-hand part of (4). The solution of the system (4) is

$$U_k = \frac{D_k}{D}. \quad (5)$$

## 5 The inverse Laplace–Carson transform and compatibility conditions

The next step of the algorithm is the inverse Laplace–Carson transform of  $U_k$ . If it is possible, than as the result we obtain the solution of the input system, which is unique (due to uniqueness of the inverse LC) and satisfies the input initial conditions. According to the character of the system – it is linear with constant coefficients – the solutions depend continuously on changing of initial conditions inside the class  $\mathbf{A}$ . That means that we found a correct (in traditional meaning) solution of the system.

The inverse Laplace–Carson transform of  $U_k$  exists under the conditions of convergence of Laplace integral of  $\widetilde{U}_k(p) = \frac{1}{p^\Gamma} U_k(p)$ , i.e. the following:

1\*  $\sigma \in \mathbf{R}^n, \sigma = (\sigma_1, \dots, \sigma_n)$ , exists such, that  $\widetilde{U}_k$  is holomorphic in the domain  $\text{Rep}_\nu > \sigma_\nu$ ;

$$2^* \lim_{p \rightarrow \infty} \widetilde{U}_k(p) = 0;$$

$$3^* \widetilde{U}_k(p) \text{ is integrable by each } p_\nu \text{ along any line } \operatorname{Re} p_\nu > \sigma_{\nu_0}, \sigma_{\nu_0} > \sigma_\nu.$$

We consider functions from the class  $\mathbf{A}$ . According to the properties of LC transform, in the fractions (5) degrees of polynomials in numerators are less than degree of the denominator, and these fractions may be represented as linear combinations of expressions of type (3). So  $2^*$  fulfills.  $3^*$  is satisfied automatically as soon as  $1^*$  takes place, due to the type (3) of expressions. So it remains to verify the implementation of  $1^*$ .

Denote a set of zeros of  $D$  by  $\mathcal{Q}$ . In the case when  $\mathcal{Q}$  has no infinite limit points at  $\mathbf{P}^+ = \{p : \operatorname{Re} p_\nu > 0, \nu = 1, \dots, n\}$  the conditions  $1^* - 3^*$  are satisfied for any initial conditions, the solution of (1) is correct.

Consider the opposite case: denote by  $\mathcal{Q}_\infty$  the subset of  $\mathcal{Q}$  with infinite limit points at  $\mathbf{P}^+$ . The condition  $1^*$  fulfills if and only if functions  $u_{\beta, \gamma}^k(x^{(\gamma)})$ ,  $k = 1, \dots, n$ , exist and are being used as initial conditions such, that  $D_k$  has zeros at  $\mathcal{Q}_\infty$  of multiplicity not less than multiplicity of corresponding zeros of  $D$ .

This demand produces requirements to the LC images of initial conditions functions, and after  $\operatorname{LC}^{-1}$  transform – to initial conditions. They turn to be dependent.

*Initial conditions, whose LC images satisfy the conditions  $1^* - 3^*$ , we call compatible.*

Thereby the following theorem is proved.

**Theorem**

*If  $\mathcal{Q}$  has no infinite limit points at  $\mathbf{P}^+$ , initial conditions, taken above, are compatible. If  $\mathcal{Q}$  has infinite limit points at  $\mathbf{P}^+$  initial conditions are compatible if and only if their images satisfy the condition:  $D_k$  has zeros at  $\mathbf{P}^+$  of multiplicity not less than multiplicity of corresponding zeros of  $D$ .*

The theorem may be considered as the condition of the existence of the inverse Laplace-Carson transform of  $U_k(p)$ .

## 6 Independence of the type of equations

As can be seen from the presented constructions, an order or a type of equations (elliptic, hyperbolic, parabolic, etc.) does not play a significant role in the method of Laplace-Carson transform. Moreover, the method allows not to reduce (or to reduce to canonical form) the problem at initial stage. However some special effects may occur in elliptic case.

## 7 Examples

In the Section 5 there was given the foundation of the method to obtain compatibility conditions, and this method was described in a general case. It would be useful not to overload the text with bulky common constructions of the implementation and to demonstrate technical details through rather transparent examples.

We present two examples: a system of the first order with composite right-hand parts and an equation of the fourth order.

*Realized in Mathematica.*

### 7.1 Example 1

To demonstrate the technics of LC algorithm let us consider in details solving a system of three equations with three unknown functions  $f(x, y, z)$ ,  $g(x, y, z)$ ,  $h(x, y, z)$  on  $\mathbf{R}_+^3$ .

$$\begin{aligned}\frac{\partial}{\partial x}f + \frac{\partial}{\partial z}g + \frac{\partial}{\partial y}h &= \phi \\ \frac{\partial}{\partial z}f + \frac{\partial}{\partial x}g + \frac{\partial}{\partial y}h &= \psi \\ \frac{\partial}{\partial y}f + \frac{\partial}{\partial x}g + \frac{\partial}{\partial z}h &= \theta\end{aligned}$$

The functions  $\phi$ ,  $\psi$ ,  $\theta$  at the right-hand parts are composite. Using Heaviside function we right them in the following way:

$$\begin{aligned}\phi &= x(H(x, y, z) - H(x - 1, y - 1, z - 1)); \\ \psi &= y(H(x, y, z) - H(x - 1, y - 1, z - 1)); \\ \theta &= z(H(x, y, z) - H(x - 1, y - 1, z - 1)).\end{aligned}$$

We shall consider the problem when values of unknown function at zeros of  $x, y, z$  are taken as initial conditions.

If we have the derivatives of the first order with respect to each variable we need nine initial conditions – three for every unknown function – at  $(0, y, z)$ ,  $(x, 0, z)$ ,  $(x, y, 0)$ , correspondingly to the order of the derivative. A requirement is a coincidence of correspondent functions values at the intersection of these planes.

Denote values of functions at these points as follows:

$$f(0, y, z) = f^x, \quad f(x, 0, z) = f^y, \quad f(x, y, 0) = f^z, \quad g(0, y, z) = g^x, \quad g(x, 0, z) = g^y, \quad g(x, y, 0) = g^z, \quad h(0, y, z) = h^x, \quad h(x, 0, z) = h^y, \quad h(x, y, 0) = h^z.$$

Denote the images of LC transform of  $f, g, h$ , respectively by  $u, v, w$ .

To be transparent in the example we denote LC images of the initial conditions functions by nine various Greek letters  $\alpha, \beta, \gamma, \delta, \tau, \varepsilon, \xi, \sigma$ :

	$(q, r)$	$(p, r)$	$(p, q)$
$f$	$\alpha$	$\eta$	$\delta$
$g$	$\varepsilon$	$\xi$	$\beta$
$h$	$\tau$	$\gamma$	$\sigma$

Table 1

In the table the first column points the functions for which the LC images of the initial conditions are considered, the first line indicates the variables upon which these images depend. Note, that in our system we have no the derivatives  $\frac{\partial}{\partial y}g$  and  $\frac{\partial}{\partial x}h$ , so we do not need  $\tau$  and  $\xi$ .

Applying the Laplace-Carson transform to the system (1) we obtain the algebraic system

$$\begin{aligned}pu + rv + qw - p\alpha - r\beta - q\gamma &= F, \\ ru + pv + qw - r\delta - p\varepsilon - q\gamma &= G, \\ qu + pv + rw - q\eta - p\varepsilon - r\sigma &= H,\end{aligned}$$

where

$$F = -e^{-p-q-r} + \frac{1}{p} - \frac{1}{p}e^{-p-q-r}; \quad G = -e^{-p-q-r} + \frac{1}{q} - \frac{1}{q}e^{-p-q-r}; \quad H = -e^{-p-q-r} + \frac{1}{r} - \frac{1}{r}e^{-p-q-r}.$$

The solution of this system is

$$\begin{aligned}
u &= -\frac{-pq^2 + pqr - pq^2r + qr^2 + pqr^2 + q^2r^2 - r^3 - qr^3}{qr(p-r)(q-r)(p+q+r)} + \\
e^{p+q+r} &\left( \frac{pq^2 - pqr - qr^2 + r^3 + p^2q^2r\alpha - p^2qr^2\alpha + pq^2r^2\beta - pqr^3\beta}{qr(p-r)(q-r)(p+q+r)} + \right. \\
&\quad \frac{-pq^2r^2\gamma + q^2r^3\gamma - pq^2r^2\delta + qr^4\delta}{qr(p-r)(q-r)(p+q+r)} + \\
&\quad \left. \frac{-pq^2r^2\epsilon + pqr^3\epsilon + pq^3r\eta - q^3r^2\eta + pq^2r^2\sigma - q^2r^3\sigma}{qr(p-r)(q-r)(p+q+r)} \right) \\
v &= \frac{-p^2q^2 - p^2q^2r + q^3r + p^2r^2 + p^2qr^2 + pq^2r^2 - qr^3 - pqr^3}{pqr(p-r)(q-r)(p+q+r)} - \\
e^{p+q+r} &\left( \frac{p^2q^2 - q^3r - p^2r^2 + qr^3 - p^2q^3r\alpha + p^2qr^3\alpha - pq^3r^2\beta}{qr(p-r)(q-r)(p+q+r)} + \right. \\
&\quad \frac{pqr^4\beta - p^2q^2r^2\gamma + pq^2r^3\gamma + pq^3r^2\delta - p^2qr^3\delta + p^3q^2r\epsilon}{qr(p-r)(q-r)(p+q+r)} + \\
&\quad \left. \frac{p^2q^3r\epsilon - p^3qr^2\epsilon - p^2q^2r^2\epsilon + p^2q^3r\eta - pq^3r^2\eta + p^2q^2r^2\sigma - pq^2r^3\sigma}{qr(p-r)(q-r)(p+q+r)} \right), \\
w &= \frac{p^2q - p^2r - q^2r - pq^2r + qr^2 + pqr^2 + q^2r^2 - qr^3}{pqr(p-r)(q-r)(p+q+r)} - \\
e^{p+q+r} &\left( \frac{-p^2q + p^2r + q^2r - qr^2 + p^2q^2r\alpha - p^2qr^2\alpha}{qr(p-r)(q-r)(p+q+r)} + \right. \\
&\quad \frac{pq^2r^2\beta - pqr^3\beta + p^2q^2r\gamma + pq^3r\gamma - pq^2r^2\gamma - q^3r^2\gamma}{qr(p-r)(q-r)(p+q+r)} + \\
&\quad \left. \frac{p^2qr^2\delta - q^2r^3\delta - pq^2r^2\epsilon + pqr^3\epsilon - p^2q^2r\eta + q^2r^3\eta - p^2qr^2\sigma + qr^4\sigma}{qr(p-r)(q-r)(p+q+r)} \right),
\end{aligned}$$

The determinant  $D$  of the system equals

$$D = -(p-r)(q-r)(p+q+r).$$

The bracket  $(p+q+r)$  is not important for solving the problem of compatibility – its zeros do not belong to  $\mathcal{Q}_\infty$ .

Consider the sets  $p=r$ ,  $q=r$ . They form the set  $\mathcal{Q}_\infty$ . We demand the numerators of the solutions be zero on these sets. To indicate that the functions of initial conditions are taken for  $p=r$  or  $q=r$  we use the notations displaced in the following table. If for a function  $p=r$  is put we use this function with the index 1, if there is put  $q=r$ , we use this function with the index 2. To demonstrate the algorithm of getting compatibility conditions display initial conditions and their transformations after substituting of points of  $\mathcal{Q}_\infty$  into the table.

	$\alpha(q, r)$	$\varepsilon(q, r)$	$\tau(q, r)$	$\theta(p, r)$	$\xi(p, r)$	$\gamma(p, r)$	$\delta(p, q)$	$\beta(p, q)$	$\sigma(p, q)$
$p=r$	$\alpha(q, r)$	$\varepsilon(q, r)$	$\tau(q, r)$	$\theta_1(r, r)$	$\xi_1(r, r)$	$\gamma_1(r, r)$	$\delta_1(r, q)$	$\beta_1(r, q)$	$\sigma_1(r, q)$
$q=r$	$\alpha_2(q, r)$	$\varepsilon_2(r, r)$	$\tau_2(r, r)$	$\theta(p, r)$	$\xi(p, r)$	$\gamma(p, r)$	$\delta_2(p, r)$	$\beta_2(p, r)$	$\sigma_2(p, r)$

Table 2

Substituting  $p = r$  and  $q = r$  into the numerators of  $u, v, w$ , we obtain a system of 6 equations, that connect functions  $\alpha, \beta, \gamma, \delta, \dots, \delta_2$ .

$$\left\{ \begin{array}{l} -q^2r + 2qr^2 - r^3 + e^{q+2r}(q^2r - 2qr^2 + r^3 + q^2r^3\alpha - qr^4\alpha + q^2r^3\beta_1 - \\ \quad qr^4\beta_1 - q^2r^3\delta_1 + qr^4\delta_1 - q^2r^3\epsilon + qr^4\epsilon) = 0 \\ q^3r - q^2r^2 - qr^3 + r^4 + e^{q+2r}(-q^3r + q^2r^2 + qr^3 - r^4 - q^3r^3\alpha + qr^5\alpha - \\ \quad q^3r^3\beta_1 + qr^5\beta_1 + q^3r^3\delta_1 - qr^5\delta_1 + q^3r^3\epsilon - qr^5\epsilon) = 0 \\ -q^2r + 2qr^2 - r^3 + e^{q+2r}(q^2r - 2qr^2 + r^3 + q^2r^3\alpha - qr^4\alpha + q^2r^3\beta_1 - \\ \quad qr^4\beta_1 - q^2r^3\delta_1 + qr^4\delta_1 - q^2r^3\epsilon + qr^4\epsilon) = 0 \\ \quad -pr^4\gamma + r^5\gamma - pr^4\delta_2 + r^5\delta_2 + pr^4\eta - \\ \quad \quad r^5\eta + pr^4\sigma_2 - r^5\sigma_2 = 0 \\ -p^2r^4\gamma + pr^5\gamma - p^2r^4\delta_2 + pr^5\delta_2 + p^2r^4\eta - \\ \quad \quad pr^5\eta + p^2r^4\sigma_2 - pr^5\sigma_2 = 0 \\ \quad p^2r^3\gamma - r^5\gamma + p^2r^3\delta_2 - r^5\delta_2 - p^2r^3\eta + \\ \quad \quad r^5\eta - p^2r^3\sigma_2 + r^5\sigma_2 = 0 \end{array} \right.$$

Solving it with respect to these variables, we get two conditions on them:

$$\begin{aligned} \alpha &= -\frac{(1-e^{-q-2r})(q-r)}{qr^2} - \beta_1 + \delta_1 + \epsilon, \\ \gamma &= -\delta_2 + \eta + \sigma_2. \end{aligned} \quad (3)$$

We may take arbitrarily all images of initial conditions except of  $\alpha$  and  $\gamma$  and obtain  $\alpha$  and  $\gamma$  according to the conditions (3).

For example, we may take the following functions in the table 1.

	$(q, r)$	$(p, r)$	$(p, q)$
$f$	$e^{-q-2r}(q-r)/(qr^2) - (q-2)/(qr^2)$	$1/(pr)$	$1/(p^2q)$
$g$	$1/(qr^2)$	$\xi$	$1/(pq)$
$h$	$\tau$	$(p-r+pr)/(p^2r^2)$	$1/(pq^2)$

Table 3

The correspondent initial conditions are the follows:

$$f^x = \frac{1}{2}((-1+2y)z^2 - (2y-z)(-2+z)H(-1+y)H(-2+z)), \quad g^x = \frac{yz^2}{2}, \quad f^y = xz, \quad h^y = \frac{1}{2}(2xz - x^2z + xz^2), \quad f^z = \frac{x^2y}{2}, \quad g^z = xy, \quad h^z = \frac{xy^2}{2}.$$

Substituting the functions  $\alpha, \beta, \gamma, \dots$  from the table 3 into the solution  $u, v, w$ , after inverse LC transform we obtain the solution of the system (1) correspondent to these initial conditions:

$$\begin{aligned} f &= \frac{1}{24} (12x^2y - 24xyz + 24yz^2 - 8z^3 - \\ &\quad H(-y+z)(8(3xy + 3y^2 - y^3 + 3xz - 3yz + 3y^2z - 3yz^2 + z^3) + \\ &\quad 8(y-z)(3x - 3y + y^2 - 2yz + z^2)H(-x+y) - \\ &\quad 12(y-z)(2+y-z)H(-1+x, -1+y) + \\ &\quad 12(y-z)(2+y-z)H(-1+x, -x+y)) - \\ &\quad 3(-2+x+z)(-2+3x-4y+3z)H(-1+y, -2+x+z) - \\ &\quad (-9x^2 + 36xy - 36y^2 - 18xz + 36yz - 9z^2)H(-1+y, x-2y+z) - \end{aligned}$$



$$\begin{aligned}
& (-12 - 24x - 9x^2 + 24y + 12xy + 6xz - 12yz + 3z^2)H(-1 - x + y, -2 - x + z) - \\
& \quad (9x^2 - 36xy + 36y^2 + 18xz - 36yz + 9z^2)H(-1 - x + y, x - 2y + z) - \\
& \quad (-12x^2 - 8x^3 + 24xz + 24x^2z - 12z^2 - 24xz^2 + 8z^3)H(-x + y, -x + z) - \\
& \quad (12 + 24y - 24z - 24yz + 12z^2)H(-1 + x, -1 + y, -1 + z) - \\
& (-12 + 24x - 9x^2 - 24y + 12xy + 24z - 18xz + 12yz - 9z^2)H(-1 + x, -1 + y, -2 + x + z) - \\
& \quad (9x^2 - 36xy + 36y^2 + 18xz - 36yz + 9z^2)H(-1 + x, -1 + y, x - 2y + z) - \\
& \quad (-24x + 9x^2 - 12xy + 24z - 6xz + 12yz - 3z^2)H(-1 + x, -x + y, -x + z) + \\
& \quad 9(x - 2y + z)^2H(-1 + x, -x + y, x - 2y + z)
\end{aligned}$$

$$\begin{aligned}
g = & \frac{1}{24} (12x^2y - 24xyz + 24yz^2 - 8z^3 - \\
& H(-y + z)(8(3xy + 3y^2 - y^3 + 3xz - 3yz + 3y^2z - 3yz^2 + z^3) + \\
& \quad 8(y - z)(3x - 3y + y^2 - 2yz + z^2)H(-x + y) - \\
& \quad 12(y - z)(2 + y - z)H(-1 + x, -1 + y) + \\
& \quad 12(y - z)(2 + y - z)H(-1 + x, -x + y)) - \\
& \quad 3(-2 + x + z)(-2 + 3x - 4y + 3z)H(-1 + y, -2 + x + z) - \\
& \quad (-9x^2 + 36xy - 36y^2 - 18xz + 36yz - 9z^2)H(-1 + y, x - 2y + z) - \\
& (-12 - 24x - 9x^2 + 24y + 12xy + 6xz - 12yz + 3z^2)H(-1 - x + y, -2 - x + z) - \\
& \quad (9x^2 - 36xy + 36y^2 + 18xz - 36yz + 9z^2)H(-1 - x + y, x - 2y + z) - \\
& \quad (-12x^2 - 8x^3 + 24xz + 24x^2z - 12z^2 - 24xz^2 + 8z^3)H(-x + y, -x + z) - \\
& \quad (12 + 24y - 24z - 24yz + 12z^2)H(-1 + x, -1 + y, -1 + z) - \\
& (-12 + 24x - 9x^2 - 24y + 12xy + 24z - 18xz + 12yz - 9z^2)H(-1 + x, -1 + y, -2 + x + z) - \\
& \quad (9x^2 - 36xy + 36y^2 + 18xz - 36yz + 9z^2)H(-1 + x, -1 + y, x - 2y + z) - \\
& \quad (-24x + 9x^2 - 12xy + 24z - 6xz + 12yz - 3z^2)H(-1 + x, -x + y, -x + z) + \\
& \quad 9(x - 2y + z)^2H(-1 + x, -x + y, x - 2y + z)
\end{aligned}$$

$$\begin{aligned}
h = & 1/24 (12xy^2 - 12x^2z - 24yz + 12xz^2 + 24yz^2 - 8z^3 + \\
& \quad 4(y - z)H(-y + z)(-2(3x - 3y + y^2 - 2yz + z^2) + \\
& \quad (2(3x - 3y + y^2 - 2yz + z^2) + 3(2 + y - z)H(-1 + x))H(-x + y) - \\
& \quad 3(2 + y - z)H(-1 + x, -1 + y)) + 12(-2 + x + z)H(-1 + y, -2 + x + z) - \\
& \quad 12y(-2 + x + z)H(-1 + y, -2 + x + z) + 9(-2 + x + z)^2H(-1 + y, -2 + x + z) - \\
& \quad 9(x - 2y + z)^2H(-1 + y, x - 2y + z) - 12(2 + x - z)H(-1 - x + y, -2 - x + z) - \\
& \quad 12x(2 + x - z)H(-1 - x + y, -2 - x + z) + 12y(2 + x - z)H(-1 - x + y, -2 - x + z) + \\
& \quad 3(2 + x - z)^2H(-1 - x + y, -2 - x + z) + 9(x - 2y + z)^2H(-1 - x + y, x - 2y + z) - \\
& \quad 12(x - z)^2H(-x + y, -x + z) - 8(x - z)^3H(-x + y, -x + z) - \\
& \quad 24(-1 + z)H(-1 + x, -1 + y, -1 + z) + 12(-1 + z)^2H(-1 + x, -1 + y, -1 + z) - \\
& \quad 12(-2 + x + z)H(-1 + x, -1 + y, -2 + x + z) + 12y(-2 + x + z)H(-1 + x, -1 + y, -2 + x + z) -
\end{aligned}$$

$$\begin{aligned}
& 9(-2+x+z)^2 H(-1+x, -1+y, -2+x+z) + 9(x-2y+z)^2 H(-1+x, -1+y, x-2y+z) + \\
& 12x(x-z)H(-1+x, -x+y, -x+z) - 3(x-z)^2 H(-1+x, -x+y, -x+z) + \\
& 24(-x+z)H(-1+x, -x+y, -x+z) + 12y(-x+z)H(-1+x, -x+y, -x+z) - \\
& 9(x-2y+z)^2 H(-1+x, -x+y, x-2y+z)
\end{aligned}$$

## 7.2 Example 2

Consider the equation of forced vibration of elastic rod:

$$\frac{\partial^2 f}{\partial x^2} + \frac{\partial^4 f}{\partial y^4} = xy.$$

Initial conditions:

$$\begin{aligned}
f(0, y) &= a(y); \quad \frac{\partial f(x, y)}{\partial x} \Big|_{x=0} = b(y); \\
f(x, 0) &= c(x); \quad \frac{\partial f(x, y)}{\partial y} \Big|_{y=0} = d(x); \\
\frac{\partial^2 f(x, y)}{\partial y^2} \Big|_{y=0} &= g(x); \quad \frac{\partial^3 f(x, y)}{\partial y^3} \Big|_{y=0} = h(x)
\end{aligned}$$

$$LC : f(x, y) \mapsto u(p, q),$$

$$\begin{aligned}
a(y) &\mapsto \alpha(q), \quad b(y) \mapsto \beta(q), \\
c(x) &\mapsto \gamma(p), \quad d(x) \mapsto \delta(p), \\
g(x) &\mapsto \sigma(p), \quad h(x) \mapsto \tau(p).
\end{aligned}$$

As a result of LC we obtain the algebraic equation:

$$p^2 u - p^2 \alpha - p\beta - qu + q\gamma qu + q\gamma = \frac{1}{pq}$$

$$D = p^2 + q^4$$

Then

$$u = \frac{1 + p^3 q \alpha + p^2 q \beta + p q^5 \gamma + p q^4 \delta + p q^3 \sigma + p q^2 \tau}{pq(p^2 + q^4)};$$

$Q$  consists of two sets of zeros of the denominator  $D = p^2 + q^4$ :

$$p = iq^2, \quad p = -iq^2.$$

At this sets the numerator of  $u$  equals correspondingly

$$\begin{aligned}
A_1 &= 1 - iq^7 \alpha - q^5 \beta + iq^7 \gamma_1 + iq^6 \delta_1 + iq^5 \sigma_1 + iq^4 \tau_1; \\
A_2 &= 1 + iq^7 \alpha - q^5 \beta - iq^7 \gamma_2 - iq^6 \delta_2 - iq^5 \sigma_2 - iq^4 \tau_2,
\end{aligned}$$

where

$$\gamma_1, \delta_1, \sigma_1, \tau_1 \text{ are the values of functions } \gamma, \delta, \sigma, \tau \text{ at } p = iq^2,$$

$$\gamma_2, \delta_2, \sigma_2, \tau_2 \text{ - at } p = -iq^2.$$

The functions with indexes 1 and 2 depend on different arguments  $iq^2$  and  $-iq^2$ , correspondingly. So it is convenient to take the originals  $c, d, g, h$  of  $\gamma, \delta, \sigma, \tau$  as data functions of initial conditions and to find  $a, b$  as compatible with them. Note that this is a characteristic speciality of equations of such type, for example of elliptic equations.

Solve

$$\begin{cases} A_1 = 0, \\ A_2 = 0 \end{cases}$$

with respect to  $\alpha, \beta$ .

Compatibility conditions on images of LC:

$$\alpha = -\frac{-q^3\gamma_1 - q^3\gamma_2 - q^2\delta_1 - q^2\delta_2 - q\sigma_1 - q\sigma_2 - \tau_1 - \tau_2}{2q^3};$$

$$\beta = \frac{i(-2i + q^7\gamma_1 - q^7\gamma_2 + q^6\delta_1 - q^6\delta_2 + q^5\sigma_1 - q^5\sigma_2 + q^4\tau_1 - q^4\tau_2)}{2q^5}.$$

Taking concrete functions  $c(t), d(t), g(t), h(t)$  of initial conditions, we obtain  $a(x)$  and  $b(x)$  as compatible with them. In such way we may define, for example, the following compatible initial conditions:

$$a = 1 - \frac{x^4}{12}, \quad b = \frac{x^5}{120}, \quad c = 1 + t^2.$$

Finally we obtain the solution satisfying the initial conditions:

$$f(t, x) = 1 + t^2 - \frac{x^4}{12} + \frac{tx^5}{120}.$$

## 8 A parallelization of solving

A parallel algorithm may be constructed on the base of the produced one.

1. The Laplace transform of right-hand parts is independent on a number of equation, it is to be produced in parallel way.
2. The Laplace transform of left-hand parts is absolutely formal and also independent, it is to be produced in parallel way.
3. Solving the algebraic system permits various methods. Most effective are based on parallel computations.
4. The inverse Laplace transform for each obtained function is to be parallelized.

## 9 Conclusion

Let us adduce advantages of the algorithm presented in the paper.

1. The Laplace–Carson transform of a system with input functions from the class **A** is being fulfilled in a symbolic way.
3. The algebraic system obtained after the Laplace transform may be solved by methods most convenient and efficient for each specific case.
  - Call a rational fraction "*a proper fraction*" if the degree of each variable (over **C**) in numerator is less than its degree in denominator.
  - Call by the class **B** a set of equations, defined by conditions
    - the solutions of algebraic system may be represented as sums of proper fractions with exponential coefficients,
    - the denominators of these proper fractions may be reduced to a product of linear functions.
4. If a system of differential equations belong to the class **B**, then the inverse Laplace transform is being fulfilled symbolically.
5. The application of Laplace–Carson transform permits to obtain compatibility conditions in a symbolic way.
6. An order of derivatives, a size of systems and for many cases types of equations are not significant for LC method.

## References

- [DahSN99] Dahiya R.S., Jabar Saberi-Nadjafi: Theorems on n-dimensional Laplace transforms and their applications. 15th Annual Conf. of Applied Math., Univ. of Central Oklahoma, Electr. Journ. of Differential Equations, Conf.02 (1999) 61-74
- [DimSp05] I.Dimovski, M.Spiridonova. Computational approach to nonlocal boundary value problems by multivariate operational calculus. Mathem. Sciences Research Journal, ISSN 1537-5978, Dec.2005, V.9, No.12, 315-329.
- [Sp09] M.Spiridonova. Operational Methods in the Environment of a Computer Algebra System. Serdica Journal of Computing, Vol. 3, No 4, (2009), 381-424.
- [NMa106] Malaschonok N.: Parallel Laplace Method with Assured Accuracy for Solutions of Differential Equations by symbolic computations. In: Computer Algebra and Scientific Computing, CASC 2006, LNCS 4196, Springer, Berlin (2006) 251-261
- [NMa110] Malaschonok N. A parallel algorithm for analytical solving of partial differential equations systems. Albanian Journal of Mathematics, V. 4, No. 4, 171-179
- [WS06] Watt S.M. Pivot-Free Block Matrix Inversion, Proc 8th International Symposium on Symbolic and Numeric Algorithms in Symbolic Computation (SYNASC), IEEE Computer Society, 2006. P. 151-155. URL: <http://www.csd.uwo.ca/watt/pub/reprints/2006-synasc-bminv.pdf>.
- [Akr89] Akritas, A.G. Elements of Computer Algebra with Applications. J. Wiley Interscience, New York (1989)
- [BH] Burghilea, D., Haller, S.: Laplace transform, dynamics and spectral geometry. arXiv:math.DG/0405037v2, 17 Jan. (2005)
- [DahSN99] Dahiya, R.S., Jabar Saberi-Nadjafi: Theorems on n-dimensional Laplace transforms and their applications. In: 15th Annual Conf. of Applied Math., Univ. of Central Oklahoma, Electr. Journ. of Differential Equations, Conf. 02 (1999) 61-74
- [DST87] Davenport, J., Siret, Y., Tournier, E.: Calcul Formel. Systemes et Algorithmes de Manipulations Algebriques. MASSON, Paris (1987)
- [GG99] Von zur Gathen, J. and Gerhard, J.: Modern Computer Algebra. Cambridge University Press (1999)
- [Mal00] Malaschonok, G.I.: Effective Matrix Methods in Commutative Domains. In: Formal Power Series and Algebraic Combinatorics, Springer, Berlin (2000) 506-517
- [Mal03] Malaschonok, G.I.: Solution of systems of linear equations by the p-adic method. Programming and Computer Software **29**(2) (2003) 59-71
- [NMa106] Malaschonok, N.: Parallel Laplace method with assured accuracy for solutions of differential equations by symbolic computations. In: Computer Algebra and Scientific Computing, CASC 2006, LNCS 4196, Springer, Berlin (2006) 251-261
- [MY97] Mizutani, N., Yamada, H.: Modified Laplace transformation method and its applications to anharmonic oscillator. 12 Feb. (1997)
- [Pod97] Podlubny, I.: The Laplace transform method for linear differential equations of the fractional order. arXiv:funct-an/9710005v1, 30 Oct. (1997)
- [PM40] Picone M.: Nuovi metodi risolutivi per i problemi d'integrazione delle equazioni lineari a derivate parziali e nuova applicazione della trasformata

- multipla di Laplace nel caso delle equazioni a coefficienti costanti. *Atti Accad.Sci.Torino*,75 (1940),1–14.
- [GD04] Goloskopov D.P. Mathematical physics equations. Solving problems in MAPLE. SPb, Piter. 2010.02.22.
- [SAS03] The method of characteristics and conversation laws. *J.of Online Math. and its App.*, url = <http://backslash/backslash/mathdl.maa.org/backslash/backslash/mathDL/backslash/4/backslash/?pa=content&sa=viewDocument&nodeId=389>, 2003.
- [BrG192] Y. Brichkov, H.-J. Glaeske, A. Prudnikov and Vu Kim Tuan ” Multidimensional Integral Transformtions” Gordon and Breach Publishers, 1992
- [DimHr09] I.H. Dimovski and V.Z. Hristov. Nonlocal Operational Calculi for Dunkl Operators. *Symmetry, Integrability and Geometry: Methods and Applications SIGMA* 5 (2009), 030, 16 pages
- [Mik56] Jan Mikusinski. *Rachunek operatorow*. Warszawa, 1953 (in Russian Jan Mikusinski. *Operatornoye ischislrniye*. M. IL. 1956)
- [Gut69] Guterman M. An operational method in partial differential equations. *SIAM J. Appl. Math.*, 17, No 2, 1969, 468-493.